



Alion[®]
TOTAL

5 l

Herbicide

A herbicide formulated as a suspension concentrate for the control of certain broadleaf and grass weeds in ratoon sugarcane.

contains:
indaziflam (Alkylazine)
Isoxaflutole (Isoxazole)





Alion®

TOTAL

SEE ENCLOSED PAMPHLET FOR FULL PARTICULARS

Reg. No. L10329 (Act No. 36 of 1947)

A herbicide formulated as a suspension concentrate for the control of certain broadleaf and grass weeds in ratoon sugarcane.

SIEN INGESLOTE PAMFLET VIR VOLLEDIGE BESONDERHEDE

Reg. Nr. L10329 (Wet Nr. 36 van 1947)

'n Onkruidodder geformuleer as 'n suspensie konsentraat vir die beheer van sekere breëblaar- en grasonkruid in oorstaanriet.

GROUP / GROEP	L ₂₉	HERBICIDE / ONKRUIDDODER
GROUP / GROEP	F ₂	HERBICIDE / ONKRUIDDODER



WARNING

HAZARD STATEMENTS:

Harmful if swallowed or inhaled. Suspected of damaging the unborn child. May cause damage to organs (Nervous system) if swallowed. May cause damage to organs (Nervous system) through prolonged or repeated exposure if swallowed. Very toxic to aquatic life with long lasting effects. Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.

PRECAUTIONARY STATEMENTS:

Do not breathe dust/ fume/ gas/ mist/ vapours/ spray.
Wear protective gloves/ protective clothing/ eye protection/ face protection.

Active ingredient:

Indaziflam (Alkylazine)..... 150 g/ℓ
Isoxaflutole (Isoxazole)..... 450 g/ℓ

Aktiewe bestanddeel:

Indaziflam (Alkylazine)..... 150 g/ℓ
Isoxaflutole (Isoxazole)..... 450 g/ℓ

REGISTERED BY / GEREGISTREER DEUR: Bayer (Pty) Ltd / (Edms) Bpk

Reg. 1968/011192/07

P.O. Box/Posbus 143, Isando, 1600 South Africa/Suid-Afrika

TEL: (OFFICE HOURS) / (KANTOORURE) (011) 921 5911

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL ASSEMBLIEF: 0861 555 777

Alion® is a registered trademark of the Bayer Group.

Alion® is 'n geregistreerde handelsmerk van die Bayer Groep.

© Bayer (Pty) Ltd

BATCH NUMBER / LOTNOMMER: See on pack

DATE OF MANUFACTURE / DATUM VERVAARDIG: Sien op houer

UN No. / VN Nr.: 3082

Bayer



RSA/0724/Alion Total 5L/
Back Booklet Code: ZA85770772D

LIFT HERE

WARNINGS:

HAZARD STATEMENTS:

- Harmful if swallowed or inhaled.
- Suspected of damaging the unborn child.
- May cause damage to organs (Nervous system) if swallowed.
- May cause damage to organs (Nervous system) through prolonged or repeated exposure if swallowed.
- Very toxic to aquatic life with long lasting effects.
- Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.

DO NOT USE PLANTS FROM TREATED SUGARCANE WITHIN 250 DAYS AFTER APPLICATION.

Store in a cool place, away from food, feed, seed or other agricultural remedies.

Keep out of reach of children, uninformed people and animals.

Re-entry - Do not enter treated area until spray deposit has dried unless wearing protective clothing.

Although **Alion Total** has been tested on most important cultivars and no significant phytotoxicity in the form of yield reduction has been recorded under normal growing conditions, this does not mean that a more sensitive cultivar might not be commercialized in the future. Where a new cultivar is encountered, large areas should not be sprayed without prior testing of **Alion Total** on that cultivar.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC AND STORAGE CONDITIONS, QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE WEEDS TO THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

PRECAUTIONARY STATEMENTS:

- Do not breathe dust/ fume/ gas/ mist/ vapours/ spray.
- Wear protective gloves/ protective clothing/ eye protection/ face protection.

Wash with soap and water after use or skin contact.

Remove soiled clothing immediately and clean thoroughly before using again.

Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and changing clothes.

Get medical advice/attention if you feel unwell.

Prevent spray drift onto other crops, grazing, rivers, dams or other areas not under treatment.

Prevent contamination of food, eating utensils, feed and drinking water.

Clean all equipment thoroughly after use and do not pollute the environment with wash water.

Do not apply where the roots of desirable plants might absorb the chemical.

Rinse the pre-mix container well.

Destroy the original container which contained **Alion Total** by puncturing and burning it. Do not use the container for any other purpose.

USE RESTRICTIONS:

- To avoid damage to follow-up crops the following waiting period should be applied between application of **Alion Total** and planting:
SUGARCANE..... 12 MONTHS
ALL OTHER CROPS..... DO NOT APPLY IN THE SEASON BEFORE REMOVAL OF THE SUGARCANE
NOTE: The abovementioned waiting periods will only be valid if the correct quantity of **Alion Total** has been applied according to the soil type and an average or above-average rainfall has occurred during and after the season in which **Alion Total** was applied. Test planting is always recommended.
- Where **Alion Total** is applied to soils which expand when wetted and crack or crumble upon drying, **Alion Total** may remain active in the soil for much longer than the above waiting periods might indicate. Therefore **Alion Total** must not be used on such soils if it is intended to plant isoxazole-sensitive crops in the foreseeable future.
- Avoid double application, e.g., overlapping spray swaths.
- Do not use on stony soil.
- Do not use on plant cane.
- Do not use on soils that contain less than 10 % clay.

RELEVANT SUBSTANCES:

Formulation inert	CAS Number	Conc. (%)
Isoxaflutole	141112-29-0	36,6
Indaziflam	950782-86-2	12,2
1,2-Benzisothiazol-3(2H)-one	2634-33-5	>0,005 – <0,05

RESISTANCE WARNING:

Alion Total contains members of the isoxazole and alkylazine group of herbicides. The mode of action of **isoxaflutole** is the inhibition of the 4-HPPD enzyme and as such it is classified by HRAC as a Group Code F₂ herbicide. Any weed population may contain individuals naturally resistant to **isoxaflutole** and other Group Code F₂ herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **isoxaflutole** or any other Group Code F₂ herbicides. The mode of action of **indaziflam** is a Cellulose biosynthesis inhibitor and is classified as a Group Code L29 herbicide by HRAC. It inhibits crystalline cellulose deposition in the plant cell wall. This action severely affects cell wall formation as well as cell elongation and division. Consequently only actively growing meristematic root and shoot tissues will be affected, hence germinating weed seeds die before emerging. Any weed population may contain or develop plants naturally resistant to **indaziflam** and other Group Code L29 herbicide. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **indaziflam** or any other Group Code L29 herbicide.

Since the presence of resistant weeds is difficult to detect prior to herbicide application, it is of the utmost importance that treated areas be inspected at regular intervals to identify the occurrence of herbicide-resistant weeds timeously. Bayer (Pty) Ltd will not accept liability for performance failures in the event of a build-up of resistant weeds resulting from inadequate resistance management practices as indicated on the label.

To delay herbicide resistance:

- Avoid exclusive, repeated use of herbicides from the same herbicide Group Code. Alternate or tank mix with products from different herbicide Group Codes.
- Integrate other control methods [chemical, cultural, biological] into weed control programmes.

For specific information on resistance management contact your local Bayer CropScience representative.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

COMPATIBILITY:

Compatibility studies have not been conducted with **Alion Total** in combination with other products.

DO NOT MIX ALION TOTAL WITH ANY OTHER PRODUCTS.

MIXING AND APPLICATION INSTRUCTIONS:

USE THE SPRAY MIXTURE WITHIN 6 HOURS AFTER MIXING

Half fill the spray tank with water. Add the pre-mix to the tank. Add the **Alion Total** to the tank at this stage and then top up with water. Keep the chemical in continuous suspension by means of agitation. After each day's spraying flush the tank out thoroughly with water. For optimum results spray as soon as possible after mixing. Accurately calibrate spray equipment prior to application. Use fan type nozzles to ensure a uniform coverage. Apply onto a clod-free surface. Use 200-300 ℓ water per ha for overall application.

APPLICATION TABLE:

CROP	SOIL CLAY PERCENTAGE	APPLICATION RATE PER HECTARE	REMARKS
SUGARCANE Ratoon cane only (Dryland and irrigated)	All soils above 10% Clay	333 mℓ Alion Total	Apply pre-emergence of the weeds.

BROADLEAF WEEDS CONTROLLED:

Ageratum conyzoides (=A. conyzoides)

Amaranthus deflexus

A. hybridus

A. spinosus

A. thunbergii

Apium leptophyllum

Bidens pilosa

Chenopodium album

C. carinatum

Conyza bonariensis (=Erigeron bonariensis)

C. sumatrensis (albida)

Convolvulus sagittatus

**Datura ferox*

**D. stramonium*

Euphorbia hirta

E. heterophylla

**Flaveria bidentis*

Gnaphalium luteo-album

G. pennsylvanicum

billy goat weed

perennial pigweed (from seed)

common pigweed

spiny pigweed

red pigweed

wild celery

common blackjack

white goosefoot

green goosefoot

horsetweed

tall fleabane

bindweed

large thorn apple

thorn apple

red milkweed

wild poinsettia

smelter's bush

jersey cudweed

Gnaphalium

BROADLEAF WEEDS CONTROLLED (cont.):

Hibiscus trionum

Hypochoeris radicata

Ipomoea obscura

**Parthenium hysterophorus*

Portulaca oleracea

Richardia brasiliensis

Schkuhria pinnata

**Senecio madagascariensis*

Sesamum triphyllum

Sida cordifolia

S. spinosa

Solanum nigrum

Tagetes minuta

bladder weed

hairy wild lettuce

small white morning glory

bastard feverfew

purslane

tropical *Richardia*

dwarf marigold

ragwort

wild sesame

heartleaf *Sida*

spiny *sida*

nightshade

khaki weed

GRASS WEEDS CONTROLLED:

Chloris virgata

Digitaria sanguinalis

D. temata

Eleusine indica

Panicum maximum

Panicum schinzii (= *P. laevifolium*)

**Paspalum dilatatum*

Rottboelia exaltata (= *R. cochinensis*)**

**Urochloa mosambicensis*

feather top chloris

crab finger-grass

black seed finger-grass

goose grass

common buffalo grass

sweet buffalo grass

common *Paspalum*

Itch grass

bushveld herringbone grass

* Later flushes of these weeds may not be adequately controlled.

** Suppression only

NOTE: Other weed species that were not present during the development trials with the product, may also be controlled to a certain degree. The registration holder does not accept any responsibility for unlisted weeds.

NOTES:

1. The spray mixtures standing for long periods of time could result either in chlorosis of the sugarcane (if sprayed onto the foliage) or in reduced residual efficacy, or both. Chlorosis symptoms are of a transitory nature and yield will not be affected significantly.
2. Application could induce chlorosis, which is transient in nature and should not affect yield.
3. Sugarcane under any form of stress is more likely to display chlorosis after **Alion Total** applications. The greater and more sustained the stress factor, the longer the plant will take to grow out of the symptoms.
4. The use of **Alion Total** on newly limed soil could cause crop chlorosis.
5. If **Alion Total** is mixed with water containing high levels of chlorine, its efficacy may be reduced.
6. Do not add any adjuvants to the spray mixture.

© Bayer (Pty) Ltd. All unauthorized reproduction of this label is forbidden.

Bayer

WAARSKUWINGS:

GEVAARVERKLARINGS:

- Skadelik indien ingesluk of ingeasem.
- Vermoede dat dit vrugbaarheid of die ongebore kind beskadig.
- Mag skade aan organe veroorsaak (senuweestelsel) indien ingesluk.
- Mag skade aan organe veroorsaak (senuweestelsel) deur langdurige of herhaalde blootstelling indien ingesluk.
- Baie giftig vir waterlewende organismes met langdurige gevolge.
- Bevat 1,2-benzisothiazolin-3-one. Mag allergiese reaksie veroorsaak.

MOET NIE PLANTE VAN BEHANDELDE RIET GEBRUIK BINNE 250 DAE NA TOEDIENING NIE.

Berg in 'n koel plek, weg van voedsel, voer, saad of enige ander landboumiddels.

Hou buite bereik van kinders, oningeligte persone en diere

Herbatedring - Moenie onbedeelde gebied binnegaan alvorens die spuitneerslag droog is nie, tensy beskermende oorklere gedra word.

Alhoewel **Alion Total** op die meeste belangrike kultivars getoets is, met geen betekenisvolle simptome van fitotoksisiteit in die vorm van oesverlagings, onder normale groeitoestande nie, beteken dit nie dat 'n meer vatbare kultivar nie in die toekomst gekommersialiseer kan word nie. Waar 'n nuwe kultivar teëgekom word, moet groot areas nie behandel word voordat **Alion Total** op daardie kultivar getoets is nie.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSEIDENHEID TOESTANDE GETOETS IS WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEINVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE GROND-, KLIMAATS- EN BERGINGSTOESTANDE, KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE ONKRUIDE TEEN DIE BETROKKE MIDDEL SOWEL AS DIE METODE, TYD EN AKKURAAHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE VAN DIE BETROKKE MIDDEL AS GEVOLG VAN DIE VERSUIM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN DIE TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMATREËLS:

VOORSORGSVERKLARINGS:

- Moet nie poeier/walm/gas/stuifsel/dampe/spuitstof inasem nie.
- Dra beskermende handskoene/beskermende klere/oogbeskerming/gesigbeskerming.

Was met seep en water na gebruik of velkontak.

Vervyder vuil klere onmiddellik en was dit deeglik voordat dit weer gebruik word.

Moenie eet, drink of rook tydens vermenging of toediening of voordat hande en gesig gewas en skoon klere aangetrek is nie.

Verky mediese advies/hulp indien siek voel.

Voorkom die oorwaai van spuitstof na ander gewasse, plantegroei, weiding, en areas wat nie onder behandeling is nie.

Verhoed besoedeling van voedsel, eetgerei, voer en drinkwater.

Maak alle apparaat deeglik skoon na gebruik en moenie die omgewing met spoelwater besoedel nie.

Moenie toedien waar die wortels van gewenste plante die middel mag opneem nie.

Spoel die voorafmeng-houer deeglik uit.

Vernietig die oorspronklike houer wat **Alion Total** bevat het deur gate daarin te maak en te verbrand. Moenie die houer vir enige ander doel gebruik nie.

GEBRUIKSBEPERKINGS:

1. Om skade aan opvolggewasse te voorkom moet die volgende wagperiodes gehandhaaf word tussen **Alion Total** toediening en plant:
SUIKERRIET..... 12 MAANDE
ALLE ANDER GEWASSE..... MOENIE TOEDIEN GEDURENDE DIE SEISOEN VOOR DIE VERWYDERING VAN DIE SUIKERRIET NIE
NOTA: Bogenoemde wagperiodes geld slegs indien die korrekte hoeveelheid **Alion Total** volgens grondsoort toegedien was en normale- of bo-gemiddelde reënval gedurende die seisoen, waarin **Alion Total** toegedien is en daarna, voorgekom het. Toetsaanplanting word altyd aanbeveel.
2. Waar **Alion Total** toegedien is op gronde wat uitsit tydens natwording en kraak of krummel tydens uitdroging soos bv. turfgronde, mag die **Alion Total** veel langer aktief bly in die grond as wat deur bostaande wagperiodes aangedui word. Gevolglik moet **Alion Total** nie op sodanige grond toegedien word indien daar beoog word om isoksasool-gevoelige gewasse in die voorsienbare toekoms daarop te plant nie.
3. Vermy dubbelbespuitings, bv., oorvleueling van spuitbane.
4. Moenie op klipperige grond gebruik nie.
5. Moenie op planriet gebruik nie.
6. Moenie gebruik op grond wat minder as 10 % klei bevat nie.

RELEVANTE STOWWE:

Formulation inert	CAS Number	Conc. (%)
Isoxaflutole	141112-29-0	36,6
Indaziflam	950782-86-2	12,2
1,2-Benzisothiazol-3(2H)-one	2634-33-5	>0,005 – <0,05

WEERSTANDSWAARSKUWING:

Alion Total insluit lede van die alkylyzine en die isoksasool-groep van onkruidodders. Die metode van werking van **isoksafutool** hierdie groep is die inhibering van die 4-HPPD ensiem en as sulks word dit deur HRAC as 'n Groepkode F₂ onkruidoder geklassifiseer. Enige populasie van 'n spesifieke onkruid mag individue insluit wat 'n natuurlike weerstand teen **isoksafutool**, of enige ander Groepkode F₂ onkruidoder het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers. Hierdie weerstandbiedende onkruid sal moontlik nie deur **isoksafutool** of enige ander Groepkode F₂ onkruidoder, beheer word nie. Die metode van werking van **indaziflam** is 'n Sellulose Biosintese Inhibeerder en is as 'n Groepkode L29 onkruidoder deur HRAC geklassifiseer. Dit inhibeer kristallyne sellulose afsetting in die plant selwand. Hierdie aksie het 'n ernstige invloed op selwand vorming sowel as verlenging en verdeling van selle. Gevolglik word slegs aktief groeiende meristematische wortel- en stamweefsel beïnvloed, vandaar sterf ontkiemende onkruid saad voor opkoms. Enige onkruid populasie mag plante insluit of ontwikkel wat 'n natuurlike weerstand teen **indaziflam** of enige ander Groepkode L29 onkruidoder het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers. Hierdie weerstandbiedende onkruid sal moontlik nie deur **indaziflam** of enige ander Groepkode L29 onkruidoder, beheer word nie.

Aangesien die voorkoms van weerstandbiedende onkruid moeilik is om te bepaal voor onkruidodertoediening plaasvind, is dit uiters noodsaaklik dat behandelde gebiede met gereelde intervalle ondersoek word om weerstand vroegtydig te identifiseer. Bayer (Edms) Bpk. sal nie verantwoordelikheid aanvaar vir enige gebrek aan prestasie in die geval van die opbou van weerstand in die onkruidpopulasie as gevolg van onvoldoende onkruidbestandheidsbestuur soos aangedui op die etiket nie.

Om weerstand teen onkruidodders te vertraag:

- Verminder die eksklusiewe, herhaaldelike gebruik van onkruidodders in dieselfde onkruidoddergroepkode. Wissel af met, of gebruik tenkmengsels van produkte in verskillende onkruidoddergroepcodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in onkruidbeheerprogramme.

Vir spesifieke inligting oor weerstandsbestuur, kontak u naaste Bayer CropScience verteenwoordiger.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHEID:

Verenigbaarheidstudies van **Alion Total** in kombinasie met ander produkte was nie gedoen nie.

MOENIE ALION TOTAL MET ANDER PRODUKTE VERMING NIE.

MENG- EN TOEDIENINGSINSTRUKSIES:

GEBRUIK DIE SPUITMENGSEL BINNE 6 UUR NA VERMENGING

Maak 'n mengsel vooraf met die aanbevole hoeveelheid **Alion Total** en 'n paar liter skoon water in 'n aparte houër. Maak die spuitnek halfvol met water. Voeg die voorafgemengde mengsel by die spuitnek. Voeg dan die ander aanbevole onkruidodder by en vul die tenk verder met water. Hou die middel in die spuitnek voortdurend in suspensie deur roering. Aan die einde van die dag se bespuiting moet die spuitnek deeglik met water uitgespoel word. Vir optimale resultate moet toediening so gou moontlik na vermenging plaasvind. Kalibreer spuittoerusting noukeurig voor toediening. Gebruik waaiertipe spuitpunte om 'n eweredige bedekking te verseker. Dien toe op die grondoppervlak wat vry is van kluite. Vir algehele toediening gebruik 200-300 l water per ha.

TOEDIENINGSTABEL:

GEWAS	GROND KLEI PERSENTASIE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
SUIKERRIET Slegs oorstaanriet (Droeland en besproeing)	Grond klei bo 10%	333 ml Alion Total	Dien voor-opkom van die onkruid toe.

BREËBLAARONKRUIDE BEHEER:

Ageratum conyzoides (= *A. conyzoides*)
Amaranthus deflexus
A. hybridus
A. spinosus
A. thunbergii
Apium leptophyllum
Bidens pilosa
Chenopodium album
C. carinatum
Conyza bonariensis (= *Erigeron bonariensis*)
C. sumatrensis (*albida*)
Convolvulus sagittatus
**Datura ferox*
**D. stramonium*
Euphorbia hirta
E. heterophylla
**Flaveria bidentis*
Gnaphalium luteo-album
G. pennsylvanicum
Hibiscus trionum
Hypochoeris radicata
Ipomoea obscura
**Parthenium hysterophorus*
Portulaca oleracea
Richardia brasiliensis
Schkuhria pinnata
**Senecio madagascariensis*
Sesamum triphyllum
Sida cordifolia
S. spinosa
Solanum nigrum
Tagetes minuta

indringer-Ageratum
meerjarige misbredie (wat van saad groei)
gewone misbredie
doringmisbredie
roomisbredie
wildeseldery
knapskérel
withondebossie
groenhondebossie
kleinskraalhans
vaalskraalhans
bobbejaantou
grootstinkblaar
stinkblaar
roomelkkruid
wilde poinsettia
smeltersbossie
roerkruid
Gnaphalium
terblansbossie
harige skaapslaai
wildepatat
Dermoinbossie
porslein
tropiese Richardia
kleinkakiebos
gifbossie
wildesesam
hartblaartaaiman
stekeltaaiman
nastergal
kakiebos

GRASONKRUIDE BEHEER:

Chloris virgata
Digitaria sanguinalis
D. temata
Eleusine indica
Panicum maximum
Panicum schinzii (= *P. laevifolium*)
**Paspalum dilatatum*
Rottboelia exaltata (= *R. cochinensis*)**
**Urochloa mosambicensis*

witpluim-Chloris
kruisvingergras
swartsaadvingergras
jongosgras
gewone buffelsgras
soet buffelsgras
gewone Paspalum
Tarentaalgras
bosveldbeesgras

* Laat stuwings van hierdie onkruid sal nie voldoende beheer word nie.

** Slegs onderdrukking

NOTA: Ander onkruidspesies wat nie tydens die ontwikkelingsproewe met die middel voorgekom het nie, kan moontlik in 'n meerdere of mindere mate beheer word. Die registrasiehouer aanvaar egter geen aanspreeklikheid ten opsigte van ongelyste onkruid nie.

NOTAS:

1. Suijtmengsels wat vir lang periodes staan mag of vergeling van die suikerriet (indien op die blare gespuj) of verkorte residuele effektiwiteit, of beide, veroorsaak.
2. Toediening mag chlorose veroorsaak, wat tydelik van aard is en behoort nie die opbrengs te beïnvloed nie.
3. Suikerriet wat onder enige vorm van stremming is sal meer geneig wees om vergeling van die blaarrandte te wys na **Alion Total** toedienings. Hoe groter en langduriger die faktor wat die stremming veroorsaak, hoe langer sal die plant neem om die simptome te ontgroei.
4. Die gebruik van **Alion Total** op pas bekalkte grond kan 'n hoë mate van gewasvergeling veroorsaak.
5. Indien **Alion Total** met water vermeng word wat hoë vlakke van chloor bevat, kan die middel se effektiwiteit verminder word.
6. Moenie bevorderingsmiddels by die spuitmengsel voeg nie.

© Bayer (Edms) Bpk. Alle ongemagtigde reproduksie van hierdie etiket word verbied.

Bayer