



Luna Flex® Fungicide

Group 7 | 3 Fungicide - Fruiting Vegetables

Luna Flex® fungicide is the newest member of the trusted Luna family of fungicides for the Eastern US. It offers long-lasting protection from key diseases in a variety of crops.

/// BENEFITS


- **Broad Spectrum**
Labeled for key crops and diseases in the Eastern US markets
- **Great Curative Control**
Enhanced curative properties to improve overall disease control
- **Long-Lasting Protection**
Extended performance over the recommended spray interval

/// LUNA FLEX® FUNGICIDE QUICK FACTS


- **Active Ingredient:** Fluopyram and Difenoconazole
- **FRAC:** Group 7 and 3 fungicide
- **Formulation Type:** Liquid, 375 SC (Soluble Concentrate)
- **Signal Word:** Caution
- **REI:** 12 hours
- **PHI:** 0 days
- **Use Rate:** 8.0 - 13.6 oz./A

Other Key Labeled Crops	Key Fruiting Vegetables Diseases
<ul style="list-style-type: none">• Apples• Bulb Vegetables• Bushberries• Citrus• Cucurbits• Pome Fruits• Stone Fruits• Strawberries	<ul style="list-style-type: none">• Anthracnose• Cercospora Leaf Spot• Early Blight• Gray Leaf Spot• Gray Mold• Powdery Mildew• Septoria Blight• Target Spot

Luna Flex® Fungicide Effectively Controls Early Blight in Tomatoes



Fresh-market tomato cv. Mountain Gem. Trial FN21USAFOX3TMD1 conducted in Waynesville, NC in 2021 by NC State University. For head-to-head comparison fungicides were applied as a full season program with seven weekly foliar applications from July 15 to September 3. Nonionic surfactant was used with all spray applications. Follow the label for rates and timings for commercial applications. The severity of early blight was evaluated weekly and were used to calculate areas under the disease progress curve (AUDPC). Means within category followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).



Trials conducted in Waynesville, NC and Vero Beach, FL in 2021 by NC State University and an independent research organization, respectively. For head-to-head comparison fungicides were applied as a full season programs with five to seven applications. Please follow the label for rates and timings for commercial applications. Nonionic surfactant was used with all spray applications. Other fungicide applications in the programs were similar across all the treatments. Luna Flex @ 8 fl oz/A and Miravis Prime @ 11.4 fl oz/A treatments were not evaluated in Waynesville, NC trial. The severity of early blight was evaluated regular intervals and data were used to calculate areas under the disease progress curve (AUDPC). Means within series followed by the same letter are not significantly different according to Fisher's Protected LSD ($P < 0.05$).

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Not all products are registered in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross and Luna Flex® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2022 Bayer Group. All rights reserved.

Luna FLEX

