The Bayer Sorghum Crop Protection Portfolio


Our portfolio of innovative solutions can help you maximize yield and performance potential from start to finish.

	Product		Group(s)†	Use Rate
Seed Treatments	Gaucho 600	Controls insects that spread viruses which reduce stand, forage production and yield potential	4 A	6.4 fl oz/100 lb of seed
Herbicides	Deg ee xtra	As soil temperature rises, patented temperature-release technology delivers just the right amount of weed control	5 15	2.0 to 3.7 qt/A
	Husl ċie °	A simple, flexible tool for control of a wide spectrum of broadleaf weeds	6 27	12.8 to 16 fl oz/A
	Huskie FX	Combines three modes of action for enhanced performance against kochia and other broadleaf weeds with built-in resistance management	4 6 27	15 to 19 fl oz/A
Herb	Roundup PowerMAX 3 HERBICIDE	Contains a unique surfactant blend with a higher consentration of glyphosate per gallon	9	20-30 fl oz/A
	RT3	Part of the Roundup Portfolio of products, RT3® herbicide is approved for pre-planting and fall burndown applications	9	32 fl oz/A
	WARRANT HERBICIDE	Acetochlor-based pre-emergence and postemergence residual herbicide with microencapsulation technology	15	1.5 to 3 qt/A
Insecticide	SIVANTO° prime	Precisely targets harmful sucking pests, such as aphids, while having minimal impact to beneficial insects	4D	7 to 14 fl oz/A*

^{*2(}ee) for control of the sugarcane aphid at lower rates in the following states: AR, KS, NE, OK, TX.

[†]Chemical mode of action Group. Please consult product labels for more information. Not all products are registered in all states and may be subject to use restrictions. Please check with local state regulatory agency for registration status.

¹For Roundup PowerMAX® 3 herbicide and RT3® herbicide applications for wheat, apply after the hard-dough stage when grain moisture is 30 percent or less. Restrictions: Do not apply more than 22 fl oz of any of these products per acre for preharvest application. Allow a minimum of 7 days between application and harvest or grazing.



² See 24(c) label for 4 to 7 fl oz/Å, 7 to 14 day PHI applications in the following states: AR, KS, NE, OK and TX. ³For Roundup PowerMAX® 3 herbicide and RT3® herbicide applications for wheat, apply after the hard-dough stage when grain moisture is 30 percent or less. Restrictions: Do not apply more than 22 fl oz of any of these products per acre for preharvest application. Allow a minimum of 7 days between application and harvest or grazing.

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information, visit www.CropScience.Bayer.us or visit us at www.AgAnytime.com to learn more about Dekalb® sorghum seed.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Technology® includes glyphosate-based herbicide technologies.

Degree Xtra® is a restricted use pesticide. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross, Degree Xtra®, Gaucho®, Huskie®, Roundup PowerMAX®, Roundup Technology®, RT 3 Powered by Roundup Technology and Design®, Sivanto® and Warrant® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-86-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2023 Bayer Group. All rights reserved. 9C0423743.