
















The Bayer

Spring Wheat Crop Protection Portfolio

Our portfolio of innovative solutions can help you maximize yield and performance potential from start to finish.

	Product	Group(s)	Use Rate		Product	Group(s)	Use Rate
Seed Treatments	 EverGol Energy	-	1 fl oz/100 lb of seed	Herbicides	 Huskie®	6 27	11 to 15 fl oz/A
	 Gaucho 600	-	0.13 to 2.4 fl oz/100 lb of seed		 Huskie COMPLETE	2 6 27	13.7 fl oz/A
	 Raxil PRO MD	-	5 to 7.5 fl oz/100 lb of seed		 Huskie FX	4 6 27	15 to 19 fl oz/A
	 Raxil PRO SHIELD	-	5 fl oz/100 lb of seed		 Olympus®	2	0.2 oz/A
Fungicides	 PROSARO®	3	6.5 to 8.2 fl oz/A		 RIMFIRE MAX	2	3 oz/A
	 PROSARO PRO	3 7	10.3 to 13.6 oz/A		 Roundup PowerMAX 3 herbicide ¹	9	20 to 30 fl oz/A
	 BAYTHROID XL	3A	1 to 2.4 fl oz/A		 RT3 ¹	9	32 fl oz/A
Insecticide					 VIOS FX	2 4	13.7 fl oz/A

¹For Roundup PowerMAX 3 herbicide, and RT3 herbicide applications for wheat, apply after the hard-dough stage when grain moisture is 30 percent or less. Restrictions: Do not apply more than 22 fl oz of any of these products per acre for preharvest application. Allow a minimum of 7 days between application and harvest or grazing.

Not all products are registered in all states. Please check with local state regulatory agency.

For more information on herbicide group numbers and site-of-action options, visit IWillTakeAction.com

Spring Wheat Application Timing

Emergence

First
Leaf

Stem
Elongations


Flag Leaf

Beginning of
Heading

Beginning of
Flowering

Ripening

Preharvest¹/
Postharvest



¹For Roundup PowerMAX 3 herbicide, and RT3 herbicide applications for wheat, apply after the hard-dough stage when grain moisture is 30 percent or less. Restrictions: Do not apply more than 22 fl oz of any of these products per acre for preharvest application. Allow a minimum of 7 days between application and harvest or grazing.

Seed Treatments Fungicides Herbicides Insecticide

For more information, visit www.CropScience.Bayer.us.

ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS. Roundup Technology® includes glyphosate-based herbicide technologies.

Baythroid® XL and Huskie® Complete are restricted use pesticides. Not all products are registered for use in all states and may be subject to use restrictions. The distribution, sale, or use of an unregistered pesticide is a violation of federal and/or state law and is strictly prohibited. Check with your local dealer or representative for the product registration status in your state. Bayer, Bayer Cross, Baythroid®, EverGol®, Gaucha®, Huskie®, Olympus®, Prosaro®, Raxil®, Rimfire®, Roundup PowerMAX®, Roundup Technology®, RT 3 Powered by Roundup Technology and Design®, Vios™ and Varro® are registered trademarks of Bayer Group. For additional product information call toll-free 1-866-99-BAYER (1-866-992-2937) or visit our website at www.BayerCropScience.us. Bayer CropScience LP, 800 North Lindbergh Boulevard, St. Louis, MO 63167. ©2023 Bayer Group. All rights reserved.